

First sighting record of Hooded Pitta *Pitta sordida* from Kaziranga National Park, India

Udayan Borthakur & Chatrapati Das

Borthakur, U., & Das, C. 2010. First sighting of Hooded Pitta *Pitta sordida* from Kaziranga National Park, India. *Indian Birds* 5 (5): 158.

Udayan Borthakur & Chatrapati Das, Aaranyak, 50 Samanway Path (Survey), Beltola, Guwahati – 781028, Assam, India.

Email: udayan@aaranyak.org

Manuscript received on 1 May 2009.

The Hooded Pitta *Pitta sordida* is found from south-western China to Philippines, and New Guinea. In India, the race *cucullata* is a summer visitor to the Himalayan foothills of Himachal to Arunachal, and north-eastern Bangladesh, and perhaps also to Meghalaya, and Nagaland. The darker *abbotti*

species for Kaziranga, 'in view of records from different parts of the Brahmanputra valley.' This is the first report of Hooded Pitta from KNP.

On 2 July 2008, we were on a field trip to the Central Range (Kohora) of KNP, as part of Aaranyak's camera-trapping team for monitoring tiger populations.

While proceeding towards Hualpath Forest Camp, between 1255–1305 hrs, we saw a pitta-like bird flying across the road (26°40.039'N 93°21.706'E). We spotted the bird again, at closer distance, in some forest undergrowth, in a patch of woodland with prominent tree species like *Machilus banbycina*, with thick undergrowth of *Calamus* sp., and creepers such as *Lea* sp. It was immediately identified as Hooded Pitta, based on its dark head, and brownish crown. We took photographs of the bird (see below), which helped confirm its identity. We observed the bird for about two minutes, before it flew away into the thickets.

References

- Barua, M. & Sharma, P. 1999. Birds of Kaziranga National Park, India. *Forktail* 15: 47-60.
- Choudhury, A. 2003. *Birds of Kaziranga National Park. A checklist*. Guwahati: Gibbon Books & The Rhino Foundation for Nature in NE India.
- Grimmett, R., Inskipp, C., & Inskipp, T. 1999. *A pocket guide to the birds of the Indian Subcontinent*. Delhi: Oxford University Press.
- Kazmierczak, K. 2000. *A field guide to the birds of the Indian subcontinent*. London: Pica Press.
- Manakadan, R., & Pittie, A. 2001. Standardized English and scientific names of the birds of the Indian Subcontinent. *Buceros* 6 (1): i-ix, 1-38.
- Rasmussen, P. C. & Anderton, J. C. 2005. *Birds of South Asia: the Ripley guide: attributes and status*. Vol 2. Washington, D.C. & Barcelona: Smithsonian Institution & Lynx Edicions.



Hooded Pitta *Pitta sordida*

is found in the Great, and Little Nicobar Islands (Rasmussen & Anderton 2005). It is categorized under 'Least Concern' in the IUCN's Red List 2008.

Kaziranga National Park (KNP) (26°35'–26°45'N 93°05'–93°40'E) is situated in the floodplain of the Brahmaputra River, in the Nowgaon, and Golaghat districts of Assam, India. A detailed checklist of birds of Kaziranga National Park, with 479 species, has listed three species of pittas, namely, Blue-naped *P. nipalensis*, Blue *P. cyanea*, and Indian *P. brachyura* (Barua & Sharma 1999). Choudhury (2003) lists the Hooded Pitta as a hypothetical