

Heronries of Tamil Nadu

S. Subramanya

PHT Scheme, 'J' Block, GKVK Campus, University of Agricultural Sciences, Bangalore 560065. Email: subbus@vnl.com

Introduction

Detailed studies on heronries in Tamil Nadu (south-eastern state in India) are very few and much of the available literature on the heronries of the region (Pittie 2005) abounds with accounts on Vedanthangal, a heronry of great antiquity (Shortt 1865, Bates 1931, Krishnan 1960, 1993, 2001, Paulraj 1985, Santharam and Menon 1991, Venkatraman and Muthukrishnan 1993, Santharam 2000, 2004, Spillett 1969, Paulraj and Kondas 1987, Baskaran 1999). The only other century-old heronry, with a recorded history, is Koonthakulam (Rhenius 1907, Webb-Pepløe 1945, Wilkinson 1961, Johnson 1971, Kumar 1980, Subramanian 2003). Of recent, information on select heronries, namely, Chitrangudi, Vettangudi, Kanjirankulam, Vettakudi-Karaivatti and a few others is available on the Internet (Anon 2005 a,b,c), in addition to occasional articles on the breeding congregation of large colonial waterbirds in the popular news media (Anon 1975, 1984a,b, MN 1980, Ratnam 1969, Jaishankar 2005, Johnson 2000, Venkataramani, 1990). The only publications that mention a few of the heronries in the state are by Krishnan (1978), and Nagulu and Rao (1983). Besides these, there have been scattered articles and notes on individual heronries (Abraham 1974, Gurusami 1988, 1994, Kumar 1980, Relton and Moses 1997, Guruswami 1999, Johnson 2000, Naganathan 2005, Karnan 2005). But for these, there has been no comprehensive work on the heronries of Tamil Nadu with an aim to understanding their status and discerning patterns in their distribution. The present paper is a step towards filling this gap in the ornithological history of the state and an effort to put the heronry scenario in the state in a proper perspective. The present work is part of a much larger nation-wide effort to understand the status and distribution of heronries in India (Subramanya 1993, 1996, 2003).

Methods

Details on the heronries of the state were collected by circulating a questionnaire to individual birdwatchers. Besides this, information available in published literature (Pittie 2005), popular print media and on the Internet was reviewed. Details of egg collections from Tamil Nadu, housed in the Oology section of The Natural History

Museum (NHM) at Tring, England (formerly known as British Museum of Natural History), were also extracted from the Internet. This information was analyzed to discern patterns in the nesting of large colonially waterbirds within the state. Details on some of the major wetlands occurring in the state were obtained from a Survey of India map of the state (scale: 1:1 million), Asian Waterfowl Census reports (van der Ven 1987, Scott and Rose 1989, Perennou et al. 1990, Perennou and Mundkur 1991, 1992) and from Vijayan et al. (2004). Location information on these wetlands was extracted from Survey of India maps. Nomenclature and systematic listing follow Manakadan and Pittie (2001).

Study Area

The present study covers the entire state of Tamil Nadu. Although, the state has been divided into 30 administrative districts, it can also be divided into three broad physiographic regions namely, the western hilly region and the plains along the coast and the north-western high elevation area. Parallel to the coast and gradually rising from it, is the broad strip of plain country in the east. It extends a little beyond the Western Ghats in Kanyakumari district and can further be sub-divided into Coromandel plains, comprising the districts of Kancheepuram, Thiruvallur, Cuddalore and Vellore; the alluvial plains of the Cauvery delta extending over Thanjavur and part of Tiruchirappalli districts and the dry southern plains in Madurai, Dindigul, Ramanathapuram, Sivagangai, Virudhunagar, Tirunelveli and Thoothukudi districts. The arid, desert-like land in the state is confined to a small area in Tirunelveli district. Tamil Nadu has a number of rivers that are relatively small and seasonal. The Cauvery delta presents some extremely distinctive physical features and is intimately associated with the life of all its broader regions. The presence of a large number of man-made wetlands (village irrigation tanks) is a characteristic feature of the lowland plains (<150 m MSL) bordering the sea (Anon 2005a).

Wetland resources of Tamil Nadu

Tamil Nadu is a water-deficient state despite c. 950 mm annual precipitation. The south-west and north-east monsoons determine

the rainfall of Tamil Nadu though some rains occur during winter (January and February) and summer (March to May). The state depends mainly on the north-east monsoon, between October and December, which is often accompanied by a depression in the south Bay of Bengal resulting in heavy rain along the coast. The Western Ghats act as a barrier, denying the state the full benefit of the south-west monsoon. Only the Nilgiris and Kanyakumari districts benefit from both monsoons (Anon 2005a).

The state has a number of seasonal rivers and the surface water resources are almost fully harnessed through 61 major reservoirs and 39,202 big and small tanks (Anon 2005a). The perennial river of the state is the Cauvery, which originates in Kodagu in neighbouring Karnataka. The Cauvery and its tributaries irrigate the fertile Coromandel plains and the Thanjavur-Nagapattinam regions, known as the granary of Tamil Nadu. The other rivers in the state are the Palar, Pennar, Vaigai and Tamiraparani. The Mettur dam forms a huge reservoir across the Cauvery, and a "Grand Anicut," a vast channel leading from it, has been constructed near Tiruchirappalli for irrigation. The waters of the Cauvery have been so well utilised that its drainage channel, the Coleroon, is always dry (Anon 2005a).

Small water storage reservoirs behind earthen dams, called tanks, dot the expanse of Tamil Nadu. Though these tanks supply many villages with drinking water, their primary purpose is irrigation. As monsoon rains fall erratically during a few months in the year, these irrigation tanks serve to store and regulate the flow of water for agricultural use, associated mainly with the production of paddy (von Oppen and Rao 1980). Paddy is the principal agricultural crop in the state, grown over nearly 2,200,000 Ha (Anon 2005a). Together, the rivers, reservoirs and the tanks provide major feeding grounds for the waterbirds in the state (Hussain and Roy 1993, Vijayan et al. 2004).

Results and discussion

Nesting species

Nineteen of the 26 species of colonially nesting large waterbirds that are known to breed in Indian heronries (Subramanya 1996), namely, Spot-billed Pelican *Pelecanus philippinus* (SBP), Little Cormorant *P. niger* (LC), Indian Shag *P. fuscicollis* (IS), Great

Cormorant *Phalacrocorax carbo* (GC), Darter *Anhinga melanogaster* (D), Little Egret *Egretta garzetta* (LE), Grey Heron *Ardea cinerea* (GH), Purple Heron *A. purpurea* (PrH), Large Egret *Casmerodius albus* (LgE), Median Egret *Mesophoyx intermedia* (ME), Cattle Egret *Bubulcus ibis* (CE), Indian Pond-Heron *Ardeola grayii* (PH), Black-crowned Night-Heron *Nycticorax nycticorax* (NH), Painted Stork *Mycteria leucocephala* (PS), Asian Openbill-Stork *Anastomus oscitans* (OBS), Glossy Ibis *Plegadis falcinellus* (GI), Oriental White Ibis *Threskiornis melanocephalus* (OWI), Black Ibis *Pseudibis papillosa* (BI), and Eurasian Spoonbill *Platalea leucorodia* (ESB), nest in Tamil Nadu (Figure 1; Appendices 1 and 2). Of these, Black-crowned Night-Heron was the commonest nesting species, occurring in about 36 existing sites (Figure 1). It was followed by Little Cormorant, Little Egret, Indian Pond-Heron and Grey Heron, with 31, 30, 29, 26 sites, respectively. Cattle Egret, Painted Stork and Spot-billed Pelican were found in less than 20 active sites each. Nests of Darter, Asian Openbill-Stork and Oriental White Ibis were found in less than 15 sites, while Great Cormorant and Eurasian Spoonbill were found to nest in 10 sites each and the rest of the species were represented in less than 10 sites (Figure 1; Appendix 1). Glossy Ibis was the least common of the nesting species, found in only two sites. Little Egret, Black-crowned Night-Heron and Little Cormorant were also the most common species in erstwhile heronries of Tamil Nadu (Figure 1, Appendix 2).

The Spot-billed Pelican appears to be surviving extremely well in Tamil Nadu with colonies reported from as many as 15 active sites. Six nesting sites of the species have been lost over the last five decades (Abraham 1974, Appendix 2). Kannan and Mankadan (2005) indicate that the Spot-billed Pelican nesting colonies may have originated from Chitrangudi and Kanjirankulam, which is not true, as both these nesting sites appear to have come into existence around the mid-1940s (Abraham 1974). In fact, Koonthakulam is the only traditional colony of the Spot-billed Pelican, which appears to have been in existence from well over a century (Webb-Peploe 1945, MN 1980), indicating that the south Indian states of Andhra Pradesh, Karnataka and Tamil Nadu have been a traditional home of Spot-billed Pelican for centuries (Subramanya 1996, 2001, 2003, Subramanya and Manu, 1996).




Figure 1. Number of nesting sites of different species in the heronries of Tamil Nadu

It is interesting to note that unlike the situation in the neighbouring state of Kerala (Subramanya 2005), the nesting sites of Great Cormorant are located inland in addition to those nesting in the reservoirs of Nilgiris district (Appendix 1). What remains of the partially submerged tree trunks within these reservoirs, serve as ideal nesting substrates for the species, a situation very peculiar to the Kerala section of the Western Ghats (Subramanya 2005).

Tamil Nadu is the only state where all the three ibis species nest together. Among these, the Oriental White Ibis is the commonest nesting species (Appendix 1). The Glossy Ibis used to be a resident in southern India (Baker and Inglis 1930) and Sri Lanka, where it was breeding well over a century ago (Henry 1971). With no records of its nesting activity thereafter, it was even considered rare in southern India (Ali and Ripley 1987, Perennou and Santharam 1990). Two nests of the species were recorded for the first time in 1993 at Koonthakulam (Sivasubramaniam *in litt.*). The Glossy Ibis is now known to nest at Vedanthangal as well (Santharam 2000). Thus, Tamil Nadu is one of the few states where Glossy Ibis has

been found to nest within Indian limits in the recent times (Subramanya 1996). The Black Ibis, usually known to nest singly or in very small loose colonies (Hume and Oates 1890, Baker 1935, Ali and Ripley 1987), seems to be largely colonial in Tamil Nadu, especially with respect to its habit of nesting along with other waterbirds in the same heronry (Appendix 1) and was recorded nesting in Tamil Nadu for the first time at Koonthakulam in 1960 (Wilkinson 1961).

The nesting colony of Grey Herons, reported near Parambikulam Reservoir in Kerala (Subramanya 2005) due to lack of proper locality details, now appears to be within the Amaravathi reserve forest, located on the shores of Amaravathi Reservoir, close to Puganodai, located some 20 km from Udumalpet, on the road to Munnar. This colony may well be the first recorded instance of the species nesting within the hills of the Western Ghats. Similarly, the egg collection data at NHM indicate that Cattle Egrets and Painted Stork, Asian Openbill-Stork and Oriental White Ibis were once nesting in the Nilgiris (Appendix 2). The Black-crowned Night-Herons, particularly those within Madurai city, are known to

frequently shift their nesting sites based on local disturbances (K. Sathasivam *in litt.*) and thus, the concentration of heronries in and around Madurai may quite possibly pertain to the same population.

Distribution of heronries

Mapping the distribution of heronries shows that not all the districts in Tamil Nadu are represented. Heronries were not reported from Tiruvannamalai, Dharmapuri, Salem, Namakkal, Karur, Pudukottai and Theni districts (Figures 2 and 3). The maximum numbers of heronries have been reported from Thirunelveli, Ramanthapuram and Kancheepuram districts, with 12 sites each, closely followed by Kanyakumari district with nine nesting sites. Six to eight heronries have been reported from each of Madurai, Chennai, and Nilgiris districts. The remaining districts hold less than four sites each. This tally is by no means an indication

of paucity or even an absence of heronries in other districts (Figure 2 and 3). There may be a large number of sites yet to be discovered and reported. For example, Pudukottai, a district with the maximum number of village irrigation tanks (Department of Economics and Statistics 2005, Vijayan et al. 2004), may indeed support a large number of heronries.

Although the present inventory of heronries in Tamil Nadu is by no means complete, looking at the three physiographic zones in Figure 5, one can distinguish six distinct clusters of heronries. Starting from the north, these clusters are distributed around Tiruvallur-Chennai-Kancheepuram; along the Cauvery River from Erode through Tiruchirapalli and the entire Cauvery delta spread over the districts of Perambalur-Nagapattinam-Tiruvarur and Thanjavur; Madurai and Sivagangai plains; Ramanthapuram districts; Tirunelveli-

Thoothukudi districts and in the extreme south around Nagercoil. The effect of the alluvial plains of the Cauvery delta, favouring paddy cultivation, on the occurrence of heronries, appears to be similar to that of Krishna and Godavari deltas in Andhra Pradesh (Subramanya 2001). When one looks at the distribution of heronries in light of the phyiographic divisions mentioned earlier, very interestingly, nearly eighty per cent of the sites are found in lower plains along the coast (Figure 4 and 5).

Figure 6 shows the distribution of all major wetlands of Tamil Nadu along with its rivers. When the distribution of these heronries are matched with the occurrence of major wetlands within Tamil Nadu (Figure 6), barring gaps of heronry information from Villupuram-Cuddalore and Tirunelveli-Virudhunagar areas, we see that they largely coincide with the distribution of major tanks, indicating the importance of these village irrigation tanks and the Cauvery delta as major foraging grounds for the species nesting in the heronries of the state.

Nesting sites

The details of nesting sites that occur in different districts along with the species nesting in them and their status is given in Appendices 1 and 2. Of the 97 sites for which information is available, 39 sites are no longer active. The rest have been active over the last 10 years. The heronries are found in varied habitats, but nearly 35 per cent of the sites areon partially submerged *Acacia nilotica* stands in village irrigation tanks. About 34 per cent of the heronries are found within human habitations while the rest are in varied situations (Table 1).

The attraction of *A. nilotica* is such that, at Koonthakulam, where birds used to nest traditionally on trees growing amidst habitation even before 1903, they have partly shifted over to nest on *A. nilotica* growing in the adjacent village tank since 1989 (Padmanabhan *in litt.*). Prior to 1989, birds used to nest only inside the village. The number of nesting species and also the number of birds visiting Koonthakulam have increased since the main breeding site has shifted to the tank close by. In selected cases where tanks with *Acacia* are found in adjoining, villages, they have been occupied by nesting birds – as seen at Peria Kollankudi-Chinna Kollankudi and Mela Selvanoor-Keela Selvanoor.




Figure 2. Number of heronries in different districts of Tamil Nadu




Figure 3. Distribution of heronries in Tamil Nadu

Key to districts: 1-Thiruvallur, 2-Chennai, 3-Vellore, 4-Kancheepuram, 5-Krishnagiri, 6-Thiruvannamalai, 7-Dharmapuri, 8-Villupuram, 9-Salem, 10-Erode, 11-The Nilgiris 12-Cuddalore, 13-Namakkal, 14-Coimbatore, 15-Perambalur, 16-Nagapattinam, 17-Karur, 18-Tiruchirapalli, 19-Thanjavur, 20-Thiruvavur, 21-Dindigul, 22-Pudukottai, 23-Theni, 24-Madurai, 25-Sivagangai, 26-Virudhunagar, 27-Ramanathapuram, 28-Tirunelveli, 29-Thoothukudi, 30-Kanyakumari.

Nesting seasons

Based on the available information, nesting seasonality of birds in heronries occurring in the state is presented in Figures 7 and 8, and Appendices 1 and 2, it is apparent that though there exists a bimodal nesting pattern among the heronries across the state. However, a majority of waterbird species considered, nest during the north-east monsoon season (Figure 8).

The heronries in western Tamil Nadu, being part of the Western Ghats, are influenced by the south-west monsoon as are the heronries of Kerala (Subramanya 2005). Only at Kudikadu and

Kachipedunatham, do the birds nest from June to October, deviating from the north-east monsoon based nesting seasons of the surrounding region. At these sites, the birds were not dependent on north-east monsoon, but on the release of water from Mettur Dam for "kuruvai" (short duration) cultivation (June-September). The birds arrived at their nesting site and began constructing nests during June and by October, the fledglings were ready to fly. This nesting activity was also known to coincide with the inflow of water to Coleroon river (Ratnam 1969). The birds at Simpson Estate heronry were recorded nesting almost throughout the

year, with certain species nesting in both the seasons (Gurusami 1994), making it to be yet another heronry that differed in its nesting periodicity from the main north-east monsoon nesting in other heronries. In Karnataka, Painted Storks nest later than Spot-billed Pelicans (Sanjay 1993), while the reverse appears to be true in Tamil Nadu. For example, at Koonthakulam, Wilkinson (1961) found that while Painted Storks had commenced nesting in January, the Spot-billed Pelicans were yet to arrive and while the pelicans were still incubating in March, the storks already had young ones in their nests.

Size and composition of nesting colonies

Several large heronries occur in Tamil Nadu. A majority of the heronries in the state are multi-species nesting sites, some of which have records of as many as 17 species nesting in them (Figure 9; Appendix 1 & 2). Nine heronries have ten or more nesting species and about 18 sites have between 5-9 species nesting in them. About 19 sites are single species colonies. The larger colony size with greater number of nesting species (Table 2) appears to be a function of age of the colony, availability of adequate nesting substrate and the quality of protection that the nesting sites enjoy (Subramanya 1996). On the other hand, the disappearance of certain nesting species over the years or decades is also known to occur as seen at Vedanthangal, where the Cattle Egret is not known to nest in recent years, while earlier studies does indicate its presence as a nesting species (Krishnan 1960, Spillett 1969, Paulraj 1985). As pointed out by Santharam (1987, 1988), such changing patterns with some of these species may largely be linked to their movement patterns within the state.

Over a century ago, Rhenius (1907) found only Spot-billed Pelican and Painted Storks breeding at Koonthakulam. In 1960, Wilkinson (1960) found Black-crowned Night-Heron and Little Egret and the Black Ibis nesting in addition, at the site. Today, especially after the birds have extended their breeding area from the village environs to nest on partially submerged *Acacia* stands in the village tank close by, Koonthakulam now boasts of 15 nesting species. Similarly, the numbers of nesting species have increased over the decades at Vedanthangal (see Santharam and Menon 1991, Santharam 2000) from the time of Shortt (1865), Packard (1903) and Bates (1931).




Figure 4. Distribution of heronries in different elevation ranges in Tamil Nadu

Traditional heronries

The popular view on the existence of heronries is that they are traditional and age-old breeding grounds for birds. This view does hold true, when one considers sites like Vedanthangal and Koonthakulam, where birds have been nesting from well over a century. Vedanthangal could be one of the oldest heronries existing in the country today. Based on documentary evidence, Bates (1931) was able to trace the existence of Vedanthangal heronry as to 1796-1798. Shortt (1865) visited Vedanthangal in March 1864, Packard (1903) in December 1902 and Bates (1931) between 1928-1930. Although documented evidences on the existence of the heronry are available since 1793, Paulraj (1985) says that its history may well date back to the Ramayana period (c. 3000 BC). Rev. Charles Theophilus Edward Rhenius, an European who lived in Tirunelveli between 1814-1838 observed the breeding of Spot-billed Pelican and Painted Storks at Koonthakulam (MN 1980) and first reported its existence in 1903. Webb-Peploe (1945) considered that the pelicanry could well have existed 60-70 years prior to 1945. In addition, sites like Simpson Estate, Chitrangudi, Vettangudi, and Kanjirankulam have existed for several decades. However, not all heronries can boast of such antiquity.

Effect of drought on nesting activity of birds

Tamil Nadu is a drought-prone state and due to vagaries of monsoon and has experienced recurrent droughts over the years (Nathan 1998). Colonially nesting waterbirds abandon breeding activities when there is lack of adequate water in their




Figure 5. Distribution of heronries in different physiographic zones in Tamil Nadu. From right to left, elevations of the three zones are 0-150m, 150-300m and 300-2000m, respectively.

feeding grounds, a majority of which are irrigation tanks. In years of poor monsoon or a monsoon failure, birds do not attempt breeding. Also, if the water in the tanks goes dry before the nesting is completed, birds are known to abandon their nests (Paulraj, 1985, Santharam 1981a, Perennou and Santharam 1991, S.S. Ramachandra Raja *in litt.*). Between 1974-1979 the entire state of Tamil Nadu experienced a severe drought and nesting did not occur in heronries of the state (Anon 1980).

Loss of heronries

Thirty-nine nesting sites have been lost over the decades in Tamil Nadu (Appendix 2). Very little information is available on the reasons for such losses. Available information indicates that disturbance by people living close to the heronries, removal of nesting substrates and poaching to be the main reasons for the loss of some of these sites (Appendix 2). Moondraidapu, an important nesting site for Spot-billed Pelicans is an example of a dramatic decline in breeding bird populations. It enjoyed protection from villagers initially (Badshah

undated), but ceased to exist by late 1980 due to poaching and felling of nesting trees. (Rajaram *in litt.*). Although people are initially tolerant towards the nesting birds, the stench that emanates from their defecation and rotting remains of fallen fish from the nests above, produces a repulsive effect on people, resulting in the eviction of nesting birds. This was also the case at Vikaas School Heronry in Madurai (T. Badarinarayanan, *verbally*), where tin cans were tied to branches within the nesting canopy to deter the birds from perching. For this very reason, the repeated attempts by birds to nest on *Acacia* stands were foiled by villagers in the tanks at Devarayaneri, Perianatham, Chamrampakkam, Arapakam, Gangaikondan, Seethaparapanallur, Nenmeni, Kaveripakkam, Thennagaram, Vaigai and Thalainayar (Badshah undated, Appendix 2). The cessation of nesting activity of birds at Simpson Estate a few years back, for reasons not yet known, is quite shocking, especially for a site which was easily the largest heronry in the state.

The felling of *Acacia* stands used by nesting birds has destroyed active heronries. Some of the tanks where the *Acacia* stands have been cut include, Velurkulam (in 2004: Robert B. Grubh, *verbally*), Periakamoi and Sakkarakottai (partially felled during 1991-1992, leaving intact only the trees on the periphery: S. Vasuki *in litt.*). At Sakkarakottai tank, a part of these peripheral trees were cut down again during 2003-2004 (S. Balachandran, *verbally*). Thus, the future of the Sakkarakottai heronry is unsure. Even the loss of heronries at Kamudi, Cheluvanoor and Pillayarkulam (Abraham 1974) may well have been due to human disturbance and the loss of nesting substrates. At Dhamal tank a vast stand of *A. nilotica*, which was a home for a large colony of cormorants and herons, was clear felled during 1969-1970 (S. T. Baskaran, *verbally*). The reasons for the loss of the small colony of Asian Openbill-Storks found by Packard (1903) or that of Miss Cockburn's Cattle Egret Colony and Walhouse's Heronry in The Nilgiris (Appendix 2) is not known.

Effect of social forestry in Tamil Nadu

The forest department in Tamil Nadu should be commended for being instrumental in the creation of a large number of heronries in the state due to the foreshore planting of village tanks with *A. nilotica*. This programme started in the early 1960s,

persisted through the 1980s and 1990s (Wilson 1979, 1986, M. Harikrishnan, *in litt.*, V. Naganathan, *verbally*). This was initiated to meet the fuel wood and small timber requirements of villagers (Wilson 1979, 1986). Under this programme, a large number of tanks were planted with *A. nilotica*. Once these *Acacia* saplings grew up, they formed a dense stand of partially submerged trees after monsoon inundation, and provided safe and ideal conditions for the nesting of colonial waterbirds. Several of these tanks with their excellent growth of *A. nilotica* support some of the large heronries in Tamil Nadu today (Table 1). Also, the most significant effect of this widespread social forestry exercise has caused a shift in the nesting pattern of large colonial waterbirds from nesting within villages to tank-based colonies. For that matter, there were very few village tanks with trees on their foreshores, which could have been used for nesting by waterbirds, prior to the commencement of social forestry programme in Tamil Nadu in early 1960s. The exception was Vedanthangal, with its 200-year history. It had over 500 partially submerged *Barringtonia* sp., trees supporting a heronry (Shortt 1865, Bates 1931, Krishnan 1960, Spillett 1969, Paulraj 1985). In those earlier times, most of the waterbirds, with a few exceptions (Table 1), appeared to have nested on trees inside villages, as seen at Koonthakulam before 1989 (Rhenius 1907, Wilkinson 1961), at Packard's colony in Chingelput (Baker 1935) and at a few others as indicated by Abraham (1974). At the same time, the commencement of heronries in tanks with *A. nilotica* stands indicates that man can indeed help in the creation and establishment of heronries. This Tamil Nadu model of foreshore planting of *A. nilotica* is worth practicing in other parts of India as well.

Importance of roost sites

As observed elsewhere in India (Subramanya 1996), many of the heronries in Tamil Nadu started as roosting sites as the basic requirements of both, the roosting, and nesting sites, of large waterbirds are almost the same. To begin with, most of these heronries start as small nascent breeding colonies with a few nesting birds. Given adequate protection, the colony grows in size, both in terms of species composition and numbers. One of the best examples of this phenomenon was the Simpson Estate heronry in Sembium, where the concerted efforts of a single individual,

turned a small roosting colony of Black-crowned Night-Herons in early 1960s, into a huge heronry with over ten nesting species and a colony size of over 20,000 birds in recent years (see Table 2, Subramanya 1996, see box). Similarly, the heronries at Udayamarthandapuram, Karikkili, Sanniyanallur and Theological College initially started as roosting sites. In certain situations, such sites have turned into safe nesting sites, mainly due to the lack of any serious disturbance and the presence of adequate protection.

Protection by villagers

Many of the heronries in Tamil Nadu are zealously protected by villagers despite the stench emanating from the nesting activities of the birds. Some of these sites include Kanjirankulam (Anon undated, Abraham 1974), Kollugudipatty (Anon 1977), Udayamarthandapuram (Anon 1990), Vettangudi (Anon 1975) and Vedanthangal (Paulraj 1985, Stairmand 1971). At Koonthakulam, any person who harmed the birds, was taken in procession around the village, with the dead bird tied around his neck (Johnson 1971), while poachers were fined with penalties at Kollugudipatty (Anon 1977). Also, when the *Acacia* stand in the tank was scheduled to be cut down at Koonthakulam, the villagers protested and prevented such a move (S. T. Basakaran, *verbally*). Even at Udayamarthandapuram, what started as an effort to curb poaching, led to the entire village passing resolutions to save the birds. An active follow-up action by local people and a village youth association finally led to the site being declared a bird sanctuary (Anon 1990). At Kollugudipatty, in an effort not to disturb nesting activity of birds, since late 1960s the entire village stopped celebrating Deepavali, the Festival of Lights, wherein bursting of noisy crackers is associated with the festivities (Anon 1977). Such initiatives by people can help to reduce human interference with nesting activity of birds. At Vettangudi, villagers are tolerant of nesting birds, and consider their arrival in large numbers as "angels of God", bringing a good harvest and various welfare programmes from the government (Karnan 2005). Similarly, a prolonged stay of birds in the village was considered as an indication of a possible double crop at Koonthakulam (Johnson 1971). These age-old notions that are connected to ground realities can be used to advantage for motivating local people and should be used in the protection

Table 1. Sites used by colonial nesting waterbirds in Tamil Nadu

Type of nesting habitat	Frequency* (%)	Name of the heronries [†]
Partially submerged trees within a tank	34.69	Dhamal Tank, Vandalur Tank, Karikkili Tank, Vedanthangal, Veeranam, Vadamugam Vellode-Periyakulam Eri, Avalpoondurai Pond, Vettakudi-Karaivetti, K. Sattanur Tank, Vadavur Tank, Kongurkulam, Udayamarthandapuram, Vettangudi Patti, Paraikanmoi, Chitrangudi, Koonthakulam, Pillayarkulam, Mela-Kelaselvanoor, Ramalingam Tank, Koonthakulam, Velurkulam, Theroorkulam, Suchindram, Manakudi, Devarayaneri, Perianatahm, Chembarambakkam, Arapakam, Gangaikondan, Seethaparapanallur, Nenmeni, Kaveripakkam, Thennagaram, Vaigai, Thalainayar
Tree on an island	2.04	Red Hill Tank, Manoli Island
Trees in and around a wetland	5.10	TV Motors, Peria Kollugudipatty, Chinna Kollugudipatty, Kanjirankulam, Eppothum Ventan
Trees close to a wetland	7.14	Kalpakkam, Thiruchinnapuram, Kudikadu, Point Calimere, Thaliyarenthal, Sankarapandiapuram, Karungulam
Trees close to reservoir	2.04	Amaravathi Reservoir, Avalanche Reservoir
Partially submerged tree trunk in a reservoir	3.06	Glenmorgan Headworks, Kamarajasagar Reservoir, Mukurthi Reservoir
Trees amidst habitation	33.67	Simpson Estate, Raja Annamalaipuram, Mylapore, Theological College, Saligramam, Packard's Chinglepat Openbill Colony, Pallipalayam, Coimbatore city, Senniyanalloor, Thatchanendal, Nagamalai, Meenakshi College Campus, Vikaasa School, Samanatham, Lady Doak College, American College, Watrup Big Tank, Sakarakottai Kanmoi, Ramanathapuram Town, Sankar Cement Campus, Poolam, Cheluvanoor, Kamudi, Ariyakulam, Koonthakulam, Thirupudai Marudur Sacred Grove, Pudugramam Osaravila, Nagercoil PWD Office, Nagercoil Nathaniel Street, Nagercoil DFO's House Compound, Nagercoil Municipal Park, Vijayanarayanam Village
Trees close to habitation	1.06	Mannampandal
Trees close to road	2.13	Battalkundu, Moondradaippu,
Trees on river bank/along a stream	3.19	Kachipedunatham, Kadavanodi, Wellington Cantonment
Tree amidst a river	1.02	Suinda Kulam
Trees close to an estuary	1.02	Adayar
Trees on seashore	1.02	Madras Crocodile Bank
Island in the gulf	1.02	Manoli
Aviary in a zoo	1.02	Arignar Anna Zoological Park

*N = 97 sites, [†] see Appendix 1 & 2 for other details

of nesting sites and to inculcate a respect for the nesting birds. There is a need for such ethos to be spread to areas where such thinking is presently absent.

Conservation of heronries

Total nesting habitat protection with availability of adequate nesting substrate would encourage an increase in the number of species and the total number of nesting birds in a heronry over time. One of the finest examples of this has been the Simpson Estate heronry where a nesting colony of about 60 birds grew over the years into a colony with nearly 20,000 birds (V. Gurusami, Viwanathan, T. Sudakar Reddy

in litt.). What has happened at Simpson Estate is an example of how man can help ensure the future survival of heronries. Vedanthangal, which has been in existence for over 200 years (Hume and Oates 1890, Baker 1935, Krishnan 1960, 1993, 2001, Paulraj 1985, Baskaran 1999), has grown in size both with respect to species and bird numbers although it suffered a setback around 35 years ago, due to the loss of the nesting substrate of *Barringtonia* trees, which were drastically reduced from over 500 trees to around 70, the reasons for which are not clear (Stairmand 1971). For that matter, a look at the species composition in heronries that existed in the past

(Appendix 2) indicates that species like Glossy Ibis and the Black Ibis have only started nesting in the state after the 1960s. This may largely be due to the opening up of additional nesting sites, when the foreshores of village irrigation tanks were planted with *A. nilotica* under the social forestry programme. Thus, the planting of *A. nilotica* has had a positive effect on the heronry scenario in the state, what with nearly forty per cent of the sites coming into existence in tanks planted with *Acacia* (Table 1).

Increasingly, heronries in Tamil Nadu are found within or close to human habitations as well (Table 1). There is a need to educate




Figure 6. Occurrence of heronries with respect to the distribution of major wetlands and rivers in Tamil Nadu

local people living close to these sites and to involve them in their conservation. Towards this end, the State Forest Department, local communities living with or close to the heronries and concerned NGOs can play a vital role in developing community-based conservation programmes. In Kokkare Bellur, Karnataka (Subramanya and Manu 1996, Manu and Jolly 2000) and Uppalapadu, Andhra Pradesh (Rao 2004), such efforts have been largely successful

Lack of protection and continued disturbance of the nesting activity of birds is catastrophic for any nesting colony, as it has been seen at Moondradaipu. It would be worthwhile if local NGOs or even individual birdwatchers in different districts of Tamil Nadu adopted heronries and worked for their long-term protection. Also,

as former roosting sites of these large waterbirds tend to grow into heronries (Subramanya 1996), all roosting sites of colonially nesting large waterbirds should be located and earmarked for protection. Protection against any form of disturbance, including poaching of eggs and birds, felling, or damage to trees utilised for nesting should be provided.

The government of Tamil Nadu has done well in protecting a considerable number of heronries. Of the 25 sanctuaries and national parks that form the protected area network in the state, 11 sanctuaries are heronries, indicating the importance given to these nesting sites. A majority of these sites have received state protection only since 1990 (Chitrangudi, Kanjirankulam, Karaivetti, Koonthankulam, Melaselvanoor-Kilaselvanoor, Udayamarthandpuram,

Vaduvvoor, Vellode), while Vedanthangal, Karikili and Vettangudi heronries have been protected prior to 1990 (Anon 2005a).

One of the factors associated with the occurrence of heronries in villages is the accumulation of copious quantities of guano beneath nesting trees. At the Kokkare Bellur pelicanry in Karnataka, every year during the nesting season, local villagers regularly spread farmyard refuse, hay, dung and several cart loads of soil or red earth below the nesting trees to collect bird droppings (Manu and Jolly 2000). At the end of the nesting season, this partially or completely decomposed guano-enriched manure is removed from below these trees and used for fertilising crops (personal observations). Mostly this is not being practiced in Tamil Nadu today, but in earlier days when birds used to nest within the Koonthakulam village in considerable numbers, harvesting of guano existed and played an important role in the protection of nesting birds (Johnson 1971). Thus, wherever possible, local people living close to the heronries should be encouraged to harvest the guano that collects below the trees. Villagers should be educated about the benefits of guano as a manure, whether it gets collected beneath the trees or falls into the water below in tank-based colonies and gets mixed with the water slated for irrigation (see Paulraj and Kondas 1987, Paulraj 1988). Development of a similar practice in the villages of Tamil Nadu harbouring heronries should encourage farmers to take active part in protecting the nesting sites. A focused campaign, effectively projecting these advantages, may help in conservation of heronries to a much greater effect.

Very recently, a heronry of Little Cormorants, Black-crowned Night-Herons and Painted Storks came into existence at the TVS Motor Company at Hosur, on the border of Tamil Nadu and Karnataka, in a wetland with trees growing in and around it, created as a part of campus planning (Subramanya et al. 2005). The management protects the birds. This site, along with the Simpson Estate heronry, remains as one of the best examples of how the corporate sector can help in creation and protection of heronries.

Future of heronries of Tamil Nadu

The information on heronries of Tamil Nadu presented in this article is by no means comprehensive. Although the present heronry survey has brought to light 97

heronries, including those that once existed, this could well be a small proportion of the real number of heronries found in Tamil Nadu. Elsewhere, intensive searches for heronries have revealed a high concentration of heronries in a single district (Subramanya 1996). The mid-winter waterfowl census estimates of large colonial nesting waterbirds (van der Ven 1987, Perennou 1989, Scott and Rose 1989, Perennou et al. 1990, Santharam 1990, Perennou and Santharam 1991, Perennou and Mundkur 1991, 1992, Rajan et al. 1992, Mundkur and Taylor 1993) show that a considerable wild population of the 19 species considered in this article, occurs in Tamil Nadu. However, the nesting populations of most of these species do not match the population estimates made during these counts. The very occurrence of a large number of village irrigation tanks, some of them very small ones, that abound in the lower plains and the area adjacent to hilly region of Tamil Nadu (Anon 2005a, Vijayan et al. 2004), and the foraging opportunities

they offer, makes one realise that there could still be many more heronries that are waiting to be discovered. Considering this, concerted efforts need to be made to develop regional inventories of heronries at district or taluk level and integrate them to project a true picture of the heronry scenario in Tamil Nadu. In this direction, individual birdwatchers, ornithologists, governmental and non-governmental agencies can contribute immensely. In addition, locating roosting and nesting sites of large colonial waterbirds should be included in the common bird programme of every birdwatcher and birdwatching NGOs in the state. In looking for heronries, it is worthwhile to explore the region around reservoirs and in regions where there are increased concentrations of waterbodies. Once an inventory has been developed, efforts should be directed to identify heronries with high density and diversity of nesting birds for long-term conservation.

Although, the availability of *Acacia* stands for nesting birds indicates an

encouraging prospect, this good omen may well be short-lived as the future of all the tank-based nesting colonies that do not enjoy state protection is uncertain. The foreshore *Acacia* plantations are eventually slated for felling after 10 years, upon which, a portion of the revenue from such felling would go to the village panchayats (Wilson 1979, 1986; Naganathan *verbally*). Although the stands that are cut down after a growth period of 10 years were to be replanted again (Naganathan *verbally*), such a move has not largely happened. Considering the importance of *Acacia* stands for nesting birds, all the tanks with nesting colonies on their *Acacia* stands should be earmarked for protection on a priority basis. NGOs with an active involvement in local community can work towards identifying priority sites and getting them declared as protected sites and educate the villager for the need to conserve them, especially by way of the manorial benefits they have to offer. Towards this, the Tamil Nadu State Forest Department should actively consider putting an end to fellings of *Acacia* stands used by birds for nesting, and all the existing sites other than those, which have already been declared as bird sanctuaries, should be accorded protection.




Figure 7. Distribution of nesting seasons in heronries across Tamil Nadu

“The Gurusami Effect”

Tamil Nadu is a state where the State Forest Department has done much for the establishment and protection of heronries. Outside this framework, there has been one individual who has devoted much of his life for conserving a heronry, the welfare of which was very dear to him. V. Gurusami's name is almost a synonym with the Simpson Estate heronry at Sembium in Chennai. What started as a small roosting site of about 60 Black-crowned Night-Herons in early 1960s, grew into one of the largest heronries in South India. Thanks to the concerted efforts of Gurusami, Tamil Nadu was able to boast of being a home for one of the finest heronries in the country. Gurusami worked as the Production Manager of Addison Paints at Simpson Estate, spread over 2.70 acres with well over 100 trees, which provided nesting space for 11 species of large colonial nesting waterbirds, numbering to over 20,000 by the turn of this century. Considering the importance of the site for a growing heronry, he rallied support of his superiors and colleagues and turned it into conservation action. His unending enthusiasm and concern for the nesting colony is noteworthy. In an era where the State Forest Departments and NGOs are working within the framework of their large mandates, Gurusami, who worked silently and relentlessly for the welfare of the nesting site, is one of the shining stars of heronry conservation in India. His unstinted efforts besides being laudable, need to be emulated everywhere in India. Gurusami has spent much of his life to set an example that amply demonstrates what a single individual can do for long-term bird conservation. The unfortunate and sudden cessation of nesting activity of birds at the estate, a few years back, must have been heart-rending for him. —S. Subramanya.

Species	Months											
	J	F	M	A	M	J	J	A	S	O	N	D
SBP												
GC												
IS												
LC												
OD												
NH												
PH												
CE												
LE												
ME												
LgE												
PrH												
GH												
PS												
OBS												
BHI												
BI												
GI												
WSB												

Figure 8. Nesting seasons of different species breeding in the heronries of Tamil Nadu. The darkest area indicates peak nesting activity.

Acknowledgements

I am very grateful to all the contributors who responded to my request for information and readily shared the details on heronries known to them. I would like to particularly mention S. Anand, Dr S. Balachandran, S. T. Baskaran, Dr Robert B. Grubb, V. Gurusami, M. Harikrishnan (former PCCF, Tamil Nadu), V. Kannan, V. Naganathan, G. Padmanabhan A. Rajaram, S. S. Ramachandra Raja, R. Ratnam, Dr A. Relton, Dr V. Santharam and late N. Suresh Babu for their kind help with information. I thank Wetland International, Malaysia, specially Dr David Lee, Dr Lalitha Vijayan and Dr. S. Narendra Prasad, Salim Ali Centre for Ornithology and Natural History, Coimbatore for providing details on some of the wetlands of Tamil Nadu; Dr. Nigel Collar for help with the egg collection data at NHM; Aasheesh Pittie for bibliographic services; K. Ratnam and Dr V. Santharam and for photocopies of articles on heronries. My study on Indian heronries was partially supported by a Small Conservation Grant from the Oriental Bird Club, U. K. Aasheesh Pittie, Gopi Sunder and V. Santharam offered comments and suggestions on earlier drafts of the manuscript.

References

Anonymous. Undated. Where birds fear neither man nor beast. *The Indian Express* (Undated).
 Anonymous. 1975. A new sanctuary for birdwatchers. *The Indian Express* 23.x.1975: 8.
 Anonymous. 1977. A village that does not celebrate Deepavali. *The Indian Express* 11.xii.1977.
 Anonymous. 1980. Migratory birds back in Nellai Dt.. *The Hindu* 13.iii.1980.
 Anonymous. 1984a. The Simpson Estate Heronry. *Newsletter for Birdwatchers* 24(5-6): 8-10.
 Anonymous. 1984b. Migratory birds arrive in Moondraidapu village. *The Hindu* 27.ii.1984.
 Anonymous. 1986. A small village but rich in migratory birds. *The Hindu* 31.x.1986.
 Anonymous. 1990. Warm welcome to winged visitors. *The Hindu* 30.iii.1990.
 Anonymous. 1998. Feathered friend love their home. *Indian Express* 21. 1.1998: 5.
 Anonymous. 2005a. State of environment report of Tamil Nadu. Internet report. <http://www.envis.tn.nic.in/> 11.v.2005.
 Anonymous. 2005b. Thiruchirappalli: Places.

URL: <http://www.all-indiatravel.com/tiruchirappalli/places-tiruchirappalli.html>. 8.iv.2005
 Anonymous. 2005c. Bird Sanctuaries in Tamil Nadu. URL: <http://www.forests.tn.nic.in/WLTBS.htm>. 11.x.2005
 Abraham, S. 1974. The Kanjirankulam breeding bird sanctuary in the Ramanad District of Tamil Nadu. *J. Bombay nat. Hist. Soc.* 70 (3): 549-552.
 Ali, S. and S. D. Ripley. 1987. *Compact handbook of the birds of India and Pakistan together with those of Bangladesh, Nepal, Bhutan and Sri Lanka*. 2nd ed. Bombay: Oxford University Press.
 Badshah, M. A. undated. Birds, animals and food. *The Indian Express*.
 Baker, E. C. S. 1935. *The nidification of birds of the Indian Empire*, Vol. 4. London: Taylor and Francis.
 Baker, H. R. and C. M. Inglis. 1930. *The birds of southern India, including Madras, Malabar, Travancore, Cochin, Coorg and Mysore*. Madras: Government Press.
 Baskaran, S. T. 1999. A threatened sanctuary. Chapter 21. In: *The dance of the Sarus: essays of a wandering naturalist*. (Ed: Baskaran, S. T.) New Delhi: Oxford University Press.
 Bates, R. S. P. 1931. *Bird life in India*. Madras: Madras Diocesan Press.
 BirdLife International. 2001. Spot-billed Pelican *Pelecanus philippensis*. In: *Threatened Birds of Asia: the BirdLife International Red Data Book*. Vol. 1. (Eds: Collar, NJ; Andreev, AV; Chan, S; Crosby, MJ; Subramanya, S; Tobias, JA) BirdLife International, Cambridge, UK, 68-103.
 Chandrasekaran, S. 1983. Ariyakulam – Abode of Painted Storks. *Bull. Madras Naturalists' Soc.* 5(1): ??.
 Daniel, P. 1980. "Koondakulam Heronry". *Hornbill* 1980 (4): 3.
 Department of Economics and Statistics. 2005. Season and crop report of Tamil Nadu 2003-04 Fasli Year – 1413. Chennai: Government of Tamil Nadu.
 Dewar, D. 1905. A list of birds found in and about Madras. *J. Bombay Nat. Hist. Soc.* 16 (3): 484-498.
 G. T. 1975. A new sanctuary for bird watchers. *The Indian Express*. 23.x.1975.
 Ganguli, U. 1964. Pelicans breeding at Moondraipu. *Newsletter for Birdwatchers* 4 (8):1-2.
 Gurusami, V. 1988. Breeding pattern of the Night Heron. *Blackbuck* 4 (2): 22-25.
 Gurusami, V. 1994. Wetland birds: Year round breeding in the Simpson Estate, Sembiam. *Blackbuck* 10 (3&4): 75-77.
 Gurusami, V. 1998. Only a breeding habitat. *Indian Express* 2.ii.1998
 Guruswami, K. 1999. The Simpson Estate wetland. *Blackbuck* 14 (3&4): 65-72.
 Henry, G. M. 1971. *The guide to the birds of Ceylon*. Bombay: Oxford University Press.
 Hume, A. O. and Oates, E. W. 1890. *Nests and eggs of Indian birds*. Vol. 3. 2nd ed. London: R.

H. Porter.

Hussain and Roy 1993. *Directory of Indian wetlands*. WWF India and AWB: New Delhi and Kuala Lumpur.

Jaishankar, C. 2005. Melaselvanoor sanctuary welcomes winged visitors. *The Hindu* 18.iv.2005.

Johnson, J. M. 1971. The heronry at Koonthakulam, Tirunelveli District, Tamil Nadu. *Newsletter for Birdwatchers*. 11 (8): 1-4.

Johnson, J. M. 2000. The birds of Koonthakulam. *Swagat* 21 (3): 50-51.

Kannan, V. and Manakadan, R. 2005. The status and distribution of Spot-billed Pelican in southern India. *Forktail* 21: 9-14.

Karnan, R. V. 2005. People's participation in the management of bird sanctuary. In: *New initiatives for bird conservation: proceedings of the national seminar on bird ecology and conservation (student supplement)*. N. S. Leela (Ed.). Bangalore: Institute for Natural Resources Conservation, Education, Research and Training.

Krishnan, M. 1960. The Vedanthangal Sanctuary for water-bird. The Madras State Forest Department.

Krishnan, M. 1978. The availability of nesting materials and nesting sites as vital factors in the gregarious breeding of Indian water-birds. *J. Bombay Nat. Hist. Soc.* 75 (Suppl.): 1143-1152.

Krishnan, M. 1993. The Vedanthangal Water-bird Sanctuary. *Humea* 1 (1): 2-8.

Krishnan, M. 2001. Vedanthangal. Chapter 57. In: *Nature's spokesman. M. Krishnan and Indian wildlife*. 1st ed. (Ed: Guha, Ramachandra). New Delhi: Oxford University Press.

Kumar, V. K. S. 1980. Koondakulam Heronry. *Hornbill* 1980 (2): 10-16.

Manakadan, R. and A. Pittie. 2001. Standardised common and scientific names of the birds of the Indian Subcontinent. *Buceros* 6 (1): 1-37.

Manu, K. and S. Jolly. 2000. *Pelicans and people: The two-tier village of Kikkare Bellur, Karnataka*. Kalpavriksh and International Institute of Environmental Development.

Menon, S. 2002. Requiem for Bird Habitats by *Madras Musings*: October 16-31, 2002: 1 & 7. Mundkur, T. and V. Taylor. 1993. *Asian waterfowl census 1993*. AWB and IWRB: Kuala Lumpur, Malaysia and, Slimbridge, U. K.

M. N. 1980. Migratory birds back in Nellai Dt. *The Indian Express* 13.iii.1980.

Naganathan, V. 2005. Bird survey report of Gulf of Munnar Biosphere Reserve-2005. Ramanathapuram, Tamil Nadu: Published by Wildlife Warden, Gulf of Munnar Marine National Park.

Nagulu, V. and J. V. R. Rao. 1983. Survey of south Indian pelicanries. *J. Bombay Nat. Hist. Soc.* 80 (1): 141-143.

Nathan, K.K. 1998. Droughts in Tamil Nadu: A qualitative and quantitative appraisal. *Drought*




Figure 9. Number of species nesting in the Heronries of Tamil Nadu

Network News 10 (3): 3-6.

Nair, M. 1996. Large Cormorant *Phalacrocorax carbo* (Shaw) breeding in the Nilgiris. *J. Bombay Nat. Hist. Soc.* 93 (1): 89.

Oppili, P. 1998. Heaven in the heart of Chennai *The Hindu* (Metro Plus). 27.iv.1998: 2.

Packard, H. N. 1903. Note on the breeding of certain herons, etc., in southern India. *J. Bombay Nat. Hist. Soc.* 15 (1): 138-139.

Paulraj, S. 1985. Studies on Vedanthangal Bird Sanctuary. Project Report-Final. Part I and II. Report submitted to the Chief Conservator of Forests, Madras 600006, Tamil Nadu.

Paulraj, S. 1988. Impact of guano deposition in Vedanthangal Water-Bird Sanctuary (Chengalpattu District, Tamil Nadu). *J. Bombay Nat. Hist. Soc.* 85: 319-324.

Paulraj, S. and Gunasekaran, G. 1988. The Vedanthangal Water-bird Sanctuary: a new breeding ground for Pelicans and Painted Storks. *J. Bombay Nat. Hist. Soc.* 85 (2): 414-415.

Paulraj, S. and S. Kondas. 1987. Two centuries of protection of a waterbird colony: how it may benefit the protectors. *Environ. Conserv.* 14: 363-364.

Paulraj, S., Manimozhi, A. and Kalyanasundaram, S. 1990. Brooding behaviour of Spot-billed or Grey Pelican (*Pelecanus philippensis*) at Arignar Anna Zoological Park, Madras, India. *Zoos' Print* V(8), 6-7.

Perennou, C. 1989. Waterfowl surveys along the Coromandel coast. *Blackbuck* 5 (4): 3-12.

Perennou, C. and V. Santharam. 1990. Status of some birds in southeastern India. *J. Bombay Nat. Hist. Soc.* 87 (2): 306-307.

Perennou, C. and V. Santharam. 1991. An ornithological survey of some wetlands in

south-east India. *J. Bombay Nat. Hist. Soc.* 87 (3): 354-363.

Perennou, C. and T. Mundkur. 1991. Asian Waterfowl Census 1991. IWRB, Slimbridge, U. K.

Perennou, C. and T. Mundkur. 1992. Asian Waterfowl Census 1992. IWRB, Slimbridge, U. K.

Perennou, C., P. Rose and C. Poole. 1990. Asian waterfowl 1990. IWRB, Slimbridge, U. K.

Perumal, I. 1967. Heronry at Samanatham. *Newsletter for Birdwatchers*. 7 (9): 7-8.

Pittie, A. 2005. *A bibliographic index to the ornithology of the Indian Subcontinent*. CD-ROM published by the author.

Rajan, V. J., T. R. Sridhar, V. S. Raghavan, P. Raman, G. Jayakumar, M. Chandrasekar, V. Gurusami, K. Vishwanathan, A. Rajaram, M. Raghuraman, B. Rajasekhar, K. V. Sudhakar, C. Ramakrishna, B. Raghunathan and R. K. G. Menon. 1992. Asian Water Fowl Counts 1992. *Blackbuck*. 8 (2): 43-47.

Ramachandran, K. 2003. Avian guests. *Hornbill* 2003(January-March): 20-21.

Rao, K. M. 2004. Report on Uppalapadu bird habitat. *Newsletter for Birdwatchers* 44 (3): 35-40.

Ratnam, K. 1969. A heronry near Lower Anicut. *The Indian Express* 20.iv.1969.

Relton, A. and A. Moses. 1997. Karavetti Bird Sanctuary – A paradise of waterfowls near Tiruchirapalli, Tamil Nadu. *J. Ecobiol.* 9 (1): 75-76.

Rhenius, C. E. 1907. Pelicans breeding in India. *J. Bombay Nat. Hist. Soc.* 17 (3): 806-807.

Sanjay, G. S. 1993. An ecological study of birds at Kikkare Bellur. Final CCV programme report submitted to WWF-India, New Delhi.

Santharam, V. 1981a. Drought in Vedanthangal. *Newsletter for Birdwatchers* 21 (5): 15.

Santharam, V. 1981b. Vedanthangal. *Newsletter for Birdwatchers* 21 (6): 7-9.

Santharam, V. 1987a. The Pond Heron – its local movements. *Newsletter for Birdwatchers*. 27(9-10): 4-6

Santharam, V. 1987b. Further notes on the local movements of the Pond Heron *Ardeola grayii*. *Newsletter for Birdwatchers*. 28(1-2): 8-9

Santharam, V. 1990. Asian waterfowl counts 1990 - a report. *Blackbuck* 6 (1): 7-11.

Santharam, V. 2000. More on Vedanthangal birds. *Newsletter for Birdwatchers* 40 (5): 57-59.

Santharam, V. 2004. More on Vedanthangal birds. *Newsletter for Birdwatchers*. 44 (1): 6-7.

Santharam, V. and R. K. G. Menon. 1991. Some observations on the water-bird populations of the Vedanthangal Bird Sanctuary. *Newsletter for Birdwatchers*. 31 (11-12): 6-8.

Scott, D. A. and P. M. Rose. 1989. Asian Waterfowl 1989. IWRB, Slimbridge, U. K..

Shortt, J. 1865. Account of a heronry, and breeding-place of other water-birds, in southern India. *J. Linn. Soc. Zool.* 8: 94-100.

Spillett, J. J. 1969. A report on wild life surveys in south and west India. November-December 1966. *J. Bombay Nat. Hist. Soc.* 65 (3): 633-

Table 2. Top ten heronries of Tamil Nadu

Name of the site [†]	No. Birds*	Species
Vedanthalangal	12,500	SBP, LC, IS, GC, D, LE, GH, GE, ME, CE, PH, NH, PS, OBS, GI, OWI, ESB
Vettangudi Patti Tank	8,000	SBP, LC, D, LE, GH, GE, ME, CE, PH, NH, PS, OBS, OWI,
Peria Kollugudipatty / Chinna Kollugudipatty	7,000	LC, LE, GH, GE, ME, CE, NH, , PS, OBS, OWI, BI, ESB
Koonthakulam	5,000	SBP, LC, IS, GC, D, LE, GE, ME, CE, NH, PS, GI, OWI, BI, ESB
Melaselvanoor-Kelaselvanoor sanctuary	4,000	SBP, LC, D, LE, GH, GE, ME, CE, PH, NH, , PS, OBS, OWI, BI, ESB
Chitragudi	4,000	SBP, LC, GC, D, LE, GH, PrH, GE, PH, NH, PS, OBS, OWI, BI, ESB
Udayamarthandapuram	3,000	LC, D, GH, ESB
Point Calimere	3,000	NH
Vadamugam Vellode-Periyakulam Eri	2,000+	LC, IS, GC, D, LE, GH, PrH, ME, CE, NH, PH, OWI, BI, ESB
Lady Doak College Heronry	2,000+	LC, LE, PH, NH

* Yearly variations are known to occur in bird numbers. † see Appendix 1 for details.

625020.

- Dr Balachandaran, S. Bombay Natural History Society, Hornbill House, Shaheed Bhagat Singh Road, Mumbai 400023.
- Dr Balasubramanian, P., Scientist, Salim Ali Centre for Ornithology and Natural History, Coimbatore 641 010 Baskaran, S.T. 9/1, 24th East Street, Tiruvanmiyur, Chennai 600041.
- Dr Iyengar, M.A.R., 17A, 2nd Avenue, Kalpakkam 603102.
- Dr Paulraj S., Former Wildlife Warden, Srivilliputtur 626125, Tamil Nadu.
- Dr Ranjit Daniels, R. J., Director, Care Earth, No.5, 21st Street, Thillaiganga Nagar, Chennai 600061.
- Dr Relton, A., Vayalur Road, Puthur, Trichirapalli 620017.
- Dr Robert B Grubb, Director, Institute For Restoration Of Natural Environment, Cosmos, 2nd Main Street, Christopher Nagar Extn., Nagercoil 629003, Kanyakumari Dist, Tamil Nadu.
- Dr Santharam, V., Institute of Bird Studies & Natural History, Rishi Valley 517352, Andhra Pradesh.
- Gurusami V., Addisons Paints & Chemicals, Simpon estate, Huzur Gardens, Sembium, Madras 600011.
- Hari Subramaniam R., 3, Chella Muthu Gownder Thottam, Sastri Nagar, Erode 638002.
- Harikrihnan, M. (IFS), Former Principal Chief Conservator of Forests, 259, Annasalai, Chennai 600006.
- Isabel Martinez and Andy Elliott, Gombu 2, 1r, 08003 Barcelona, Catalunya, Spain.
- Kannan R. (IAS), Former Collector, Periyar District, Erode 638011.
- Karthikeyan, M., First year M.Sc., Wildlife Biology, AVC College, Mannampandal, 609305.
- Kavin D. Paulraj, s/o Dr. P. Dayanandan, Madras Christian College, Tambaram, Chennai 600059.
- Krishnan L.V., No.6, 11th Street, Kalpakkam 603102.
- Kumaran Sathasivam, 29, Jadamuni Koil Street, Madurai 625001.
- Manoj V. Nair, #34, Thoppil Nagar, Kumarapuram, Trivandrum 695011, Kerala.
- Menon R.K.G., 18, IV Main Road, Gandhi Nagar, Adayar, Chennai 600020.
- Mohammed Ali S., Wildlife Preservation Society, 247, Main Road, Mettupalayam 641301.
- Mrs. Chukerbuti I.E., Bonnie Nestle, Gorkha Hill, Wellington, Nilgiris, Tamil Nadu.
- Padmanabhan G., Porunai Nature Club, B-64, SPIC Nagar, Thoothukudi 628005.
- Palani V.(I.F.S.), Former District Forest Officer, Kanyakumari District.
- Palaniappan, N.S. (I.A.S.), Former Collector, Periyar District, Erode 638011.
- Paramasivan, G., Honorary Chairman, Institute of Nature Studies, The Sengunthar Higher Secondary School, Erode 638001.
- Raajalingaraja, S.V., Arignar Anna Zoological park, Vandalur, Chennai 600048.
- Raghunatha V., D-211, Staff Quarters, GKVK Campus, University of Agricultural Sciences, Bangalore 560065.
- Rajaram A., C2, #29, 4th Seaward Road, Valmiki Nagar, Thiruvanmiyur, Chennai 600041.
- Ramachandra Raja S.S., Vice-President, Wildlife Society of Ramanad District, 58, P.S.K. Nagar, Rajapalayam 626108.
- Ramaswamy, G., Lecturer in Wildlife Biology, AVC College, Mannampandal, 609305.
- Sanjay G.S., 2208/B, Sadhana Road, K.R. Extension, Tumkur 572101, Karnataka.

- 663.
- Stairmand D. A. 1971. Vedanthalangal. *Newsletter for Birdwatchers* 11 (7): 6-9.
- Subramanian, K. S. 2003. Koonthakulam Bird Sanctuary: An overview. *Zoos' Print* XVIII: 27-28.
- Subramanya, S. 1993. Information Request: Catalogue of Indian Heronries. *IWRB News*. 9:11.
- Subramanya, S. 1996. Distribution, status and conservation of Indian heronries. *J. Bombay Nat. Hist. Soc.* 93 (3): 459-486.
- Subramanya, S. 2001. Heronries of Andhra Pradesh. *Mayura* 13: 1-27.
- Subramanya, S. 2003. Distribution, status and conservation of Indian heronries. In: (Daniel, J. C., and Gayatri W. Ugra., Eds.) *Petronia: Fifty years of post-independence ornithology in India (A centenary dedication to Dr. Salim Ali)*. 1st ed. Bombay: Bombay Natural History Society and Oxford University Press.
- Subramanya, S. 2005. Heronries of Kerala. *Malabar Trogon* 3 (1): 2-15.
- Subramanya, S., M. B. Krishna, S. Karthikeyan, V. Cavale and O. C. Naveein. 2005. *Observations and recommendations for the Painted Stork colony at the TVS Motor Company premises*. Unpublished report submitted to TVS Motor Company, Hosur, Tamil Nadu.
- Subramanya, S. and Manu, K. 1996. Saving the Spotbilled Pelican: A successful experiment. *Hornbill*: 1996(2): 2-6.
- van der Ven, J. 1987. Asian Waterfowl 1987: Midwinter bird observations in some Asian countries. Slimbridge, England: IWRB.
- Venkataramani, G. 1990. Annual haunt for migratory birds. *The Hindu* 28.xii.1990.
- Venkatraman, C and Muthukrishnan, S.1993.

- Density of waterbirds at Vedanthalangal Bird Sanctuary, T.Nadu. in: Bird Conservation Strategies for the nineties and beyond. (Eds.): Abraham Verghes, S. Sridhar and A.K. Chakravarthy. OSI, Bangalore. Pp. 55-60.
- Vijayan, V. S., S. N. Prasad, L. Vijayan, and Muralidharan, S. (2004): Inland Wetlands of India: Conservation Priorities. Salim Ali Centre for Ornithology and Natural History. Coimbatore.
- von Oppen, M. and K. V. S. Rao. 1980. Tank irrigation in semi-arid tropical India. Part 1: Historical development and spatial distribution. Andhra Pradesh, India: ICRISAT.
- Webb-Peploe, C. G. 1945. Notes on a few birds from south of the Tinnevely district. *J. Bombay Nat.Hist. Soc.* 45: 425-426.
- Wilkinson, M. E. 1961. Pelicanry at Kundakulam, Tirunelveli District. *J. Bombay Nat. Hist. Soc.* 58 (2): 514-515.
- Wilson, J. 1979. Social forestry in Tamil Nadu. *Indian Forester* 105: 700-706.
- Wilson, J. 1986. Management of community forests in Tamil Nadu. *Indian Forester* 112: 305-313.
- Zari, A. A. and A. R. Rahmani. 2005. Annotated avifauna of the Upper Nilgiris, Western Ghats, Tamil Nadu, India. *Buceros*: 10 (1): 1-46.

List of contributors

- Ambudoss Arvind, 7/99, NGGO Colony, Nagamalai, Madurai 625 019.
- Anand Naik K.G., Former District Forest Officer, Periyar District, Erode 638 003.
- Anand, S., Education Officer, WWF-India. Tamil Nadu State Office, 297, TTK Road, Alwarpet, Chennai 600018.
- Dr Badarinarayanan, T., 262, II Main Road, Gomathipuram, Sivagangai Road, Madurai

Sankaran, K., #14, V.N. Nagar, Tiruchirapalli 620002.
 Shivaramamurthy, Simpon estate, Huzur Gardens, Sembium, Madras 600011.
 Sivaperuman, C., 1 year M.Sc., Wildlife Biology, AVC College, Mannampandal, 609305.
 Sivasubramaniam K., Asst Prof.(Seed Tech.) Tamil Nadu Agricultural University, Coimbatore 641003.
 Sudakar Reddy T., Addisons Paints & Chemicals, Simpon estate, Huzur Gardens, Sembium, Madras 600011.
 Sugato Dutt (IFS), Former Wildlife Warden, No. 50, IV Main Road, Gandhinagar, Adayar, Chennai 600020.

Susanthkumar, C., 'Prakruthi', Indiranagar, Peroorkada P.O., Trivandrum 695 005.Suresh Babu N., Canara Bank, Vanross Jn., Trivandrum 695039.
 Vallanattan, A., Kandavarayanpatti 623 203, Pasumpon Thevar Thirumaganur District.
 Vasuki S., Former Lab Assistant c/o Wildlife Warden, Gulf of Mannar, Marine National Park, 76/1, Ramanatahapuram.
 Velayudan V.S., 24, Banker' Colony, Kumaran Nagar, Vayaloor Road, Tiruchirapalli 620017.
 Viswanathan K., Addisons Paints & Chemicals, Simpon estate, Huzur Gardens, Sembium, Madras 600011.

S. Subramanya is a Ph.D. in agricultural ornithology (University of Agricultural Sciences, Bangalore, Karnataka). He is a Member: Heron Specialist Group, IUCN, Switzerland and a Trustee of New Ornis Foundation. He has co-edited *Threatened Birds of Asia: the BirdLife International Red Data Book* (2001; BirdLife International: Cambridge).

Appendix 1. Details of existing heronries in Tamil Nadu

District	Name	Nesting Spp.	Season	Source
Kancheepuram	Arignar Anna Zoological Park	SBP,LE,PS	Nov-March	S.V. Raajalingaraja, Paulraj et al. 1990
	Vandalur Tank	LC	July	S.V. Raajalingaraja
	Karikkili	SBP,LC,ME,CE,PH,NH, PS,OBS	Oct-March	Anon 2005 a & c, S. Dutt, A. Rajaram, Krishnan (1978),
	Edayur	SBP+	Oct-March	Kannan and Manakadan 2005
	Vedanthalangal	SBP,LC,IS,GC, D,LE,GH,LgEME, CE,PH,NH,PS,OBS,GI,OWI, ESB	Oct-March	S. Dutt, Kevin D. Paulraj, R.K.G. Menon, Shortt 865, Bates 1931, Krishnan 1960, 1993, 2001, Spillett 1969, Siromoney 1984, Santharam 1981b, Paulraj 1985, Paulraj and Kondas 1987, Paulraj and Gunasekaran 1988, Santharam and Menon 1991, Venkatraman and Muthukrishnan 1993, Baskaran 1999, Santharam 2000, 2004, BirdLife International. 2001, Anon 2005 a, c,
	Kalpakkam	LC,D,LE,GH,CE,NH, PS	Oct-Jan	Rajaram A..L.V. Krishnan, M.A.R. Iyengar
Dharmapuri	TV Motors	LC,NH,PS	Feb-June	S. Subramanya, Subramanya et al, 2005
Erode	Pallipalayam	LC IS,D,LE,CE,PH,NH	May-June	S. Mohamad Ali,
	Vadamugam Vellode-Periyakulam Eri	LC,IS,GC,D,LE,GH,PrH,ME, CE,PH,NH, OWI,BI,ESB	Nov-Jan	R.Kannan, K.G. Anand Naik, G. Paramasivan, R. Hari Subramaniam, Anon 2005a
	Avalpoondurai Pond	LC,GC,LE,CE,PH,NH,BI	Nov	G. Paramasivan
The Nilgiris	Glenmorgan Headworks	GC	Sept	Nair 1996
	Kamarajasagar Reservoir	GC	Sept	Nair 1996.
	Mukurthi Reservoir	GC	Sept-Oct	S. Subramanya
	Wellington Cantonment	PH	May	I.E. Chukerbuti
	Avalanche Reservoir	GC	July	Zarri and Rahmani 2005
Cuddalore	Thiruchinnapuram	LE,GH,LgE,PH,OBS	Nov-March	S. Anand
	Veeranam	LE,GH,NH	Nov-March	S. Anand
Coimbatore	Coimbatore city	NH	Oct	S. Mohamad Ali,
	Amaravathi Reservoir	GH,NH	July-Oct	Isabel Martinez and Andy Elliott
Perambalur	Kudikadu	LC,LE,GH,ME,PH,NH	June-Oct	Ratnam 1969
	Kachipedunatham	LC,LE,GH,ME,PH,NH	June-Oct	Rathnam 1969
Nagapattinam	Mannampandal	LE,CE,PH,NH	Nov-Dec	G.Ramaswamy, C. Sivaperuman and M.Karthikeyan.
	Point Calimere	NH	Oct-Dec.	S. Anand
Tiruchirapalli	Vettakudi-Karaivetti	SBP,LC,D,LE,GH,NH,PS,OBS, OWI,ESB	Nov-March	A.Relton, BirdLife International. 2001,Anon 2005a,b
	K. Sattanur Tank	CE,PH,NH	Nov-March	V.S. Velayudan

District	Name	Nesting Spp.	Season	Source
Thiruvavur	Vadavur Tank	LE,GH,PH	Oct-March	Anon 2005a, c
	Udayamarthandapuram	LC,D,GH,ESB	Aug-Jan	K. Sivasubramanian, Anon 1990, Anon 2005a,c
Dindigul	Kongurkulam	LE,CE,PH,NH	Oct-Jan	S.T.Baskaran
	Battalkundu (Kodaikanal foothills)	PH,NH	Jan-Dec	A.Relton
Madurai	Nagamalai	PH	Oct-Dec.	Ambudass Arvind,
	American College	LC, LE,PH,NH	Nov-Jan	T. Badarinarayanan
Sivagangai	Vettangudi Patti Tank	SBP,LC, D,LE,GH,LgE,ME,CE, PH,NH,OBS,PS,OWI	Oct-April	V. Naganathan, A. Rajaram, G.S.Sanjay and Ragunatha, V, G.T. 1975, Naganathan 2005a,c
	Peria Kollugudipatty/ Periyakollankudi	LC,LE,GH,LgE,ME,NH,CE, PS,OBS,OWI,BI,ESB	Oct-April	A. Vallanattan, G.T. 1975, Anon 1977
	Chinna Kollugudipatty/ Chinnakollankudi	LC,LE,GH,LgE,ME,CE,NH, PS,OBS,OWI,BI,ESB	Oct-April	A. Vallanattan, G.T. 1975, Anon 1977
Virudhunagar	Watrup Big Tank	SBP,LC,OD,NH	Dec-March	S. Balachandran, Kannan and Manakadan 2005
	Sankarapandiapuram	LE,GH,PS,BI	Feb-May	S. Paulraj
Ramanathapuram	Parai Kanmoi	SBP,LC,D,LE,GH,PrH,PH,NH, PS,OBS,OWI	Nov-March	S. Balachandran, S. Vasuki
	Kanjirankulam	SBP,LC,LE,GH,NH,PH, PS,OWI,BI	Oct-March	S.Vasuki, Abraham 1974, BirdLife International. 2001, Anon 2005 a,c
	Sakarakottai Konamai	SBP,LC,LE,GH	Nov-March	S. Vasuki
	Ramanathapuram town	LC,LE,NH,PH	Oct-March	A.Relton
	Chitrangudi	SBP,GC,LC,D,LE,GH,PrH,LgE, PH,NH,PS,OBS,OWI,BI,ESB	Oct-July	P. Balasubramanian, S.S. Ramachandra Raja, Sanjay, G.S. and V. Raghunatha, S.Vasuki, BirdLife International. 2001, Anon 2005a,c
	Thaliyarethal Tank	GH,LE	Feb-March	Naganathan 2005
	Manoli Island	GH	Jan-March	Naganathan 2005
	Mela-Keelaselvanoor sanctuary	SBP,LC,D, LE,GH,LgE,ME,CE, PH,NH,PS,OBS,BI,OWI,ESB	Oct-April	V. Naganathan, Anon 2005a, Jaishankar 2005, Naganathan 2005
Thoothukudi	Eppothum Ventan	LC,LE,NH,PS,OBS	Oct-March	Robert B. Grubb,
Tirunelveli	Sankar Cement Campus	LC,GH,PH	Nov-Dec	A.Relton
	Poolam	SBP,PS	Jan	Kannan and Manakadan 2005
	Ariyakulam	SBP,PH,NH,PS	Jan-July	G. Padmanabhan, A. Rajaram, Robert B. Grubb, Sanjay, G.S. and V. Raghunatha, Siva Subramanian, C. Susanthakumar, Chandrasekaran 1983, Nagulu and Rao 1983, Anon 1984, BirdLife International. 2001
	Koonthakulam	SBP,GC,IS,LC,D,LE,NH,CE,LgE, ME, PS,GI, OWI,BI, ESB	Nov/Feb-May/ June	Rhenius 1907, Webb-Peploe 1945, Wilkinson 1961, Johnson 1971, Kumar 1980, BirdLife International. 2001, Subramanian 2003, Anon 2005a
	Thirupudai Marudur Sacred Grove	LC,NH,PS	March-June	Robert B. Grubb
Kanyakumari	Pudugramam Osaravila	CE	July-August	C. Susanthakumar
	Nagercoil DFO's House Compound	LC,LE,NH	April-July	Robert B Grubb
	Nagercoil Nathaniel Street	LC,LE,NH	April-July	Robert B Grubb
	Nagercoil PWD Office	PH,NH	April-July	Sureshbabu, N.,
	Nagercoil Municipal Park	PH,NH	April-July	Sureshbabu, N.
	Suinda Kulam	LC,LE,GH,PrH,NH	April-July	Sureshbabu, N.
	Suchindram Tank	LC,LE,GH,PrH,NH,OBS	NK	V.Palani
	Manakudi Tank	LC,LE,NH,GH,PrH,OBS	NK	V.Palani

*Abbreviations as per text, NK= Not known

Appendix 2. Details of erstwhile heronries of Tamil Nadu

District	Name of the heronry [†]	Nesting species*	Season	Last seen	Reasons for loss	Source
Thiruvallur	Red Hill Tank	NH	Aug-Dec	early 1960		Dewar 1905
Chennai	Simpson Estate	IS, LC, D, NH,	June-Sept; PH, CE, LE, ME, LgE	2002 Oct-March	NK	V. Gurusami, Shivarama- murthy, K.Viwanathan, T.Sudakar Reddy
	Theological College	LC, IS, LE, PH, NH	Oct-March	1998		V. Santharam, Anon 1998, Gurusami 1998, Menon 2002, Oppili 1998
	Saligramam	NH	July/August.	1982		V. Santharam
	Adayar	LE, CE, PH, NH	Oct-March	pre-1978		Krishnan 1978
	Raja Annamalaipuram	LE, CE, PH, NH	Oct-March			Krishnan 1978
	Mylapore	LE, CE, PH, NH	Oct-March			Krishnan 1978
Vellore	Kaveripakkam	NK	Oct-March	pre-1970	Disturbance by people	Badshah undated
Kancheepuram	Chamrampakkam	NK	Oct-March			
	Packard's Openbill Colony	OBS	NK	pre-1935	NK	Baker 1935
	Madras Crocodile Bank	LE, LgE, ME, CE, PH, NH ,	Oct-March	pre-1993		R. J. Ranjit Daniels
	Dhamal Tank	LC, LE, GH, NH, OBS	Oct-March	1969-70	Felling of Acacia trees	Krishnan 1978, S.T.Baskaran, A. Rajaram
	Perianatham	NK	Oct-March	pre-1970	Disturbance by people	Badshah undated
Arapakam	NK	Oct-March				
Villupuram	Thennagramam	NK	Oct-March			
Nilgiris	Miss Cockburn's Colony	CE	NK	1864	NK	BNHM Egg collection
	Walhouse's Colony	PS, OBS,OWI	NK	pre-1861		Anon 1986
Nagapattinam	Senniyanalloor	LC, LE, PH, OBS	Oct-Dec.	pre-1986		
Tiruchirapalli	Devarayaneri	NK	Oct-March	pre-1970	Disturbance by people	K.Sankaran, Badshah undated
Thanjavur	Thalainayar	NK	Oct-March			Badshah undated
Madurai	Thatchanendal	GH	Oct-March	early 1990s	NK	T.Badarinarayanan
	Vikaasa School	LC, LE, NH,	Nov-Jan	mid 1990s	Birds evicted	Kumaran Sathasivam; T. Badarinarayan
	Meenakshi College campus	NH	Nov-Jan		Disturbance by people	Kumaran Sathasivam; T. Badarinarayan
	Vaigai	NK	Oct-March	pre-1970		Badshah undated
	Samanatham	LE, GH	Oct-April	1967	NK	Perumal (1967):
	Lady Doak' College	NH	Oct-April	2004-2005	Birds evicted	S.T. Bhaskaran, T. Badarinarayanan
Ramanathapuram	Cheluvanoor	SBP, PS	Oct-April	pre-1950s	Disturbance by people	Abraham 1974
	Nenmeni	NK	Oct-March	pre-1970		Badshah undated
	Kamudi	SBP, PS	Oct-March	pre 1950s		Abraham 1974
	Pillayarkulam	SBP, PS	Oct-March			Abraham 1974
	Gangaikondan	NK	Oct-March	pre 1970	Disturbance by people	Badshah undated
	Seethaparanallur	NK	Oct-March			Badshah undated
	Ramalingam Tank	SBP, LC, LE, PS	Jan-March	pre 1961	NK	Ganguli 1964
	Moondradaippu	SBP, LC, LE, PS	Jan-March- June	1988	Poaching and felling of trees	A. Rajaram, R.J. Ranjit Daniels, Ganguli 1964, MN 1980, Anon 1984, Bird Life International. 2001
	Tirunelveli					
	Velurkulam	LC, NH, PS	March-June?	2002	Felling of Acacia trees	Robert B Grubh
	Karungulam	D, LE, CE, NH	NK	pre 1978	NK	Krihnanan 1978
	Vijayanarayanam Village	SBP, LC, PS	March-June	pre 1980		MN 1980
Thoothukudi	Kadavanodi	LC	Jan-April	1972	Poaching of nestlings	Daniel (1980)
Kanyakumari	Theroorkulam	LC, LE, GH, PrH, NH, OBS	NK	1993	Felling of Acacia trees	V.Palani